UNITED EQUITY Item # 10001 CALF STARTER

18% TEXTURIZED CALF STARTER MON 37.5

MEDICATED Calf Starter for the prevention coccidiosis caused by Eimeria bovis and Eimeria zuernii (EXCLUDING VEAL CALVES)

INGREDIENTS

Grain products, plant protein products, molasses products, processed grain by-products, calcium carbonate, vegetable oil, salt, pork fat (preserved with BHA), calcium phosphate, magnesium oxide, hydrolyzed yeast, sodium carboxymethyl-cellulose, vitamin E supplement, ferrous sulfate, manganese proteinate, zinc proteinate, roughage products, Yucca schidigera plant extract, selenium yeast, copper proteinate, calcium pantothenate, niacin supplement, cobalt proteinate, mineral oil, vitamin A supplement, vitamin D-3 supplement, menadione nicotinamide bisulfite (source of vitamin K activity), riboflavin supplement, vitamin B-12 supplement, pyridoxine hydrochloride, choline chloride, folic acid, thiamine mononitrate, ethylenediamine dihydroiodide, and biotin.

ACTIVE DRUG INGREDIENTS: Monensin, 37.5 g/ton AVAILABLE IN 50 lb (22.69 kg) BAGS or BULK.

FEEDING DIRECTIONS

Each pound of United Equity 18% Texturized Calf Starter MON 37.5. contains 18.75 milligrams of Monensin. Feed United Equity 18% Texturized Calf Starter MON 37.5 continuously at the rate of 0.75 to 5.33 pounds per 100 pounds of bodyweight per day to provide 0.14 to 1.0 milligrams Monensin per pound of bodyweight per day up to a maximum of 200 milligrams Monensin per head per day. Introduce calf starter at 4 days of age and keep fresh to encourage consumption. Offer clean, fresh water constantly. Avoid feeding hay prior to weaning. Wean calves at 4 - 6 weeks of age when eating a minimum of 3 pounds per day of United Equity 18% Texturized Calf Starter MON 37.5 for two consecutive days. Following this desired consumption, gradually transition calves to a grower ration during the next 2 to 4 weeks.

- Supplement pellet is full TRT (Total Replacement Technology)
 which makes the minerals more readily available to digest.
 - Contains Bio-Mos[®], a yeast-based prebiotic. It helps to build natural defenses by supporting a healthy gut microbiome.
 - Contains De-Odorase® to help control ammonia levels, manure odors and other harmful gasses associated with animal waste decomposition. It helps create a healthier environment for animals and their caretakers.
 - Also contains Monovet[®] for the prevention of coccidiosis.
- Made with molasses and CLE to help get a freshly weaned calf a good start on feed.
- Local-sourced corn and heavy test weight oats for ample nutrition.
- Depending on milk consumption, feed intake, and desired wean weight, feed 1 to 2 bags of UE Calf Starter per calf.

GUARANTEED ANALYSIS	
Crude Protein, min	18.0%
Crude Fat, min	3.1 %
Crude Fiber, max	8.5 %
Acid Detergent Fiber, max	9.5 %
Calcium, min	0.7 %
Calcium, max	1.2 %
Phosphorus, min	0.4 %
Salt, min	0.2 %
Salt, max	0.7 %
Selenium, min	0.28 ppm
Vitamin A, min	8,600 IU/lb

WARNINGS

Do not use in calves to be processed for veal. A withdrawal period has not been established for this product in pre-ruminating calves. CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Do not feed undiluted. Monensin medicated cattle and goat feeds are safe for cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats.

This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for dairy and is not intended for other species. Caution: Follow label directions. Feeding this feed containing added selenium at levels higher than recommended is not permitted.

